



Effect of Dietary Supplementation of *Albizia lebbek* Seed Oil on the Fatty Acid Composition of Weaner Rabbits

Alagbe J. O.

Department of Animal Production and Nutrition, Sumitra Research Farm, Gujarat, India

Email: demsonfarms@yahoo.com

Article History

Received: July 15, 2020

Revised: August 8, 2020

Accepted: August 17, 2020

Published: August 20, 2020

Copyright © 2020 ARPG & Author

This work is licensed under the Creative Commons Attribution International

 CC BY: Creative Commons Attribution License 4.0

Abstract

An experiment was carried out to determine the effects of dietary supplementation of *Albizia lebbek* seed oil (ALO) on the fatty acid composition of weaner rabbits. 50 weaned rabbits of mixed breed and sexes, aged between 6-7 weeks with an average initial body weight of 460 ± 1.3 were randomly assigned into five dietary treatments of ten rabbits per group; each group was further divided into 5 replicates consisting of two rabbits each. Basal diet was formulated to meet the nutritional requirements of rabbits according to National Research Council [1]. Rabbits in treatment 1, 2, 3, 4 and 5 were supplemented with ALO at 0 %, 0.1 %, 0.2 %, 0.3 % and 0.4 % respectively. Feed and water was given ad libitum and the experiment lasted for 12 weeks. The results showed that significant differences ($P < 0.05$) were observed in saturated fatty acid (SFA), polyunsaturated fatty acid (PUFA) and omega-6/omega-3 ratio (n-6: n-3) values obtained. Rabbits fed diet containing 0.4 % ALO had the highest PUFA value (54.17 %), followed by T4 (53.01 %), T3 (45.13 %), T2 (37.61 %) and T1 (26.93 %) respectively. Similarly (n-6: n-3) composition in T5 (3.65 %) increased in T5 fed 0.4 % ALO compared with T1 (1.38 %) fed 0% ALO. Anthrogenic index were significantly ($P < 0.05$) different among the treatments. It can be concluded that supplementation of ALO at 0.4 % highly influenced the composition of fatty acid in rabbit meat.

Keywords: Rabbit; *Albizia lebbek* seed oil; Essential oil; Fatty acid.

1. Introduction

The use of essential oil as feed additives in livestock production is becoming an interest since the ban on the use of antibiotics by the European Union in 2006, since it alleviates the problem of antimicrobial resistance, residues in animal products and the danger posed to human health.

Essential oils (EOs) are mixture of fragrant, safe and volatile compounds, which are usually originated from plant and are named with the aromatic characteristics considering the origin of plant [2]. It has also been reported to contain several bioactive or phytochemicals which confer it the ability to perform multiple biological activities such as: antimicrobial, antiviral, antifungal, antiparasitic and antidiarrheal because of the presence of tannins, flavonoids, saponins, phenols and alkaloids [3, 4]. Phytochemicals are also known phytochemicals are natural bioactive compounds that are derived from plants and incorporated into animal feed to enhance productivity [5]. The antimicrobial and modulatory properties of EOs are recent focus for agricultural applications because of a desire on the part of livestock farmers to reduce the use of unnatural chemicals in feed production [6].

EOs consists of two compounds, the terpenes and phenylpropenes, their concentration vary according to the plant, parts of the plant, geographical origin, season and processing techniques [5, 7]. They also play a role in appetite stimulation, improvement of enzyme secretion related to food digestion and immune activation [8] especially oil obtained from *Albizia lebbek*.

Albizia lebbek is a large deciduous tree belonging to the family Leguminosae (Fabaceae). The plant is found in West Africa and some parts of Asia including India. According to Everist [9] *Albizia lebbek* tree range from medium to large of multi-stemmed spreading habit. The leaves are traditionally used for feeding ruminant animals and are also used for the treatment of asthma, malaria, conjunctivitis and diarrhea Mohammed, *et al.* [4] and its seeds are easily extracted by hand from the pods by hand or by crushing the pods and winnowing to obtain the oil Mohammed, *et al.* [4] and also good sources of carbohydrates, proteins, fat soluble vitamins minerals and fatty acids. The seed oil is abundant in oleic and linoleic acid at 5.3 % and 78.5 % respectively [10]. *Albizia lebbek* seed contains alkaloids, saponins, tannins and flavonoids which have a high therapeutic value [11].

Previous research has proven that changes in the lipid composition of animal diet can alter the composition of meat fat [12]. Alagbe, *et al.* [13], reported that modification of fatty acid via feeding provided the best results in

monogastric animals. So many essential oil have been experimented in animals, for instance, *Centella asiatica* oil [14], linseed oil [15], flaxseed oil [16], garlic oil [17], thymol and cinnamaldehyde oil [18] and most recently *Albizia lebbbeck* oil. The use of *Albizia lebbbeck* essential oil will bridge a gap between food safety and livestock production.

Therefore this experiment was designed to determine the effect of dietary supplementation of *Albizia lebbbeck* seed oil on the fatty acid composition of weaner rabbits.

2. Materials and Methods

2.1. Site of the Experiment

The experiment was carried out at Division of Animal Nutrition, Sumitra Research Farm, Gujarat, India during the month of December to February, 2020.

2.2. Collection of Test Material and Preparation

Healthy seeds of *Albizia lebbbeck* were harvested within the farm premises in India, it was later authenticated at the Herbarium unit of Department of Biological Sciences on the farm and assigned a voucher specimen number MM- 1256 AL. The seeds were separated manually from their seed coats and sundried for one (1) week. The dried seeds were granulated into coarse particles using a grinder in the laboratory. *Albizia lebbbeck* oil (ALO) was extracted from the coarse particles using soxhlet extraction method. It was later poured into air tight labelled container and subjected to further analysis.

2.3. Experimental Animals and Their Management

Fifty (50) weaned rabbits of mixed breed and sexes, aged between 6-7 weeks with an average initial body weight of 460 ± 1.3 were obtained from a commercial farm in India and used in the experiment. Before the commencement of the experiments, animals were given prophylactic treatment of Ivermectin injection at the rate of 0.2 ml/rabbit administered subcutaneously and broad spectrum antibiotics (Oxytrox L. A[®]), multivitamins (Biovit super[®]) were given intramuscularly at the rate of 0.2 ml and 0.1ml/ rabbit respectively. The rabbits were housed individually in a special all wired cages of dimension 60cm × 60 cm (length and width) equipped with concentrate drinkers and feeders. Five dietary treatments of ten rabbits per group were used for this experiment. The animals were allowed one-week adjustment period during which they were fed with basal diet. Feed and water were given *ad libitum* by 7:00 am and 17:00 pm and all management practices will be strictly observed throughout the experimental period which will last for 12 weeks.

2.4. Experimental Set-Up

Experimental diet was formulated to meet the nutritional requirements of rabbits according to [National Research Council \[1\]](#), as presented in [Table 1](#)

Treatment 1: Basal diet + 0.0 % ALO

Treatment 2: Basal diet + 0.1 % ALO

Treatment 3: Basal diet + 0.2 % ALO

Treatment 4: Basal diet + 0.3 % ALO

Treatment 5: Basal diet + 0.4 % ALO

2.5. Measurements

Feed intake was recorded daily and body weight gain was recorded weekly, feed conversion ratio was calculated by dividing the total feed intake by weight gain. Mortality was also recorded as it occurs and the experiment lasted for 12 weeks.

2.6. Fatty Acid Analysis

At the end of the experiment, five rabbits were randomly selected from each treatment for fatty acid analysis (FA). Meat (longissimus dorsi muscles) lipids from freeze dried, grounded samples were extracted with chloroform-methanol (2:1 v/v; [Folch, et al. \[19\]](#) with slight modification as described by [Elshater, et al. \[20\]](#). After extraction, FAs in the residual fat were esterified, using acid and base catalyzed methods as described by [Elshater, et al. \[20\]](#). Fatty acid methyl esters (FAMES) analysis was performed by gas chromatography mass spectrometry (GC-MS; Mussek-QM- 2010 plus, China) equipped with electron impact (EI) detector. Separations of FAs were carried out on capillary column Model 7009 A, Punjab Technologies, India (30m × 0.32 mm × 0.25 μm) using helium as carrier gas. Column temperature was held at 50°C for 1 min, and then the temperature was raised up to 150°C at the rate of 15°C per min. Temperature was later increased to 175°C at the rate of 2.50°C and hold for 5 min and finally increased to 220°C at the rate of 2.50°C per min and kept for 5 min. The identification of the peaks was made by comparison of the equivalent chain length with those of authentic fatty acid methyl esters. Peak areas were determined automatically using the Agilent gas chromatography chemstation software. The fatty acid concentrations are expressed in percentage of the sum of total identified peaks measured in each sample.

2.7. Chemical Analysis

Phytochemical analysis was carried out on the test material (ALO) using standard methods [21]. Percentage composition of flavonoids, saponin, phytate, alkaloids, tannin and oxalate were carried out according to procedures outlined by Harbone [22] and Boham and Kocipai [23].

2.8. Statistical Analysis

All data collected was subjected to one-way analysis of variance (ANOVA) using SPSS (23.0) and significant means will be separated using Duncan multiple range tests [24] significant will be declared if $P \leq 0.05$.

Table-1. Ingredient composition of experimental diets

Ingredient	Quantity
Maize	36.00
Wheat offal	25.00
Palm kernel meal	20.20
Groundnut cake	10.00
Soya meal	5.10
Limestone	1.00
Bone meal	2.00
Lysine	0.10
Methionine	0.10
*Premix	0.25
Salt	0.25
Calculated analysis (% DM)	
Crude protein	16.44
Crude fibre	8.82
Ether extract	2.46
Calcium	1.34
Phosphorus	0.50
Energy (Kcal /kg)	2500.8

* Premix supplied per kg diet :- Vit A, 7,000 I.U; Vit E, 8mg; Vit D3, 3000I.U, Vit K, 3mg; Vit B2, 5.5mg; Niacin, 25mg ; Vit B12, 16mg ; Choline chloride, 120mg ; Mn, 5.2mg ; Zn, 25mg ; Cu, 2.6g ; Folic acid, 2mg ; Fe, 5g ; Pantothenic acid, 10mg ; Biotin, 30.5g ; Antioxidant, 56mg

Table-2. Phytochemical analysis of *Albizia lebbek* seed

Parameters	% Composition	*Safe recommended level
Tannins	2.11	11.20
Saponin	6.33	7.02
Alkaloids	0.11	0.55
Oxalate	0.12	1.30
Glycosides	0.10	0.50

Table-3. Effect of *Albizia lebbek* seed oil on the fatty acids profile of rabbit meat

Fatty acids	T1	T2	T3	T4	T5	SEM
C12:0	3.88 ^a	2.00 ^b	1.95 ^b	1.91 ^b	1.82 ^b	0.01
C14:0	3.22 ^a	2.62 ^b	2.40 ^b	2.29 ^b	2.00 ^c	0.18
C16:0	22.9 ^a	17.3 ^b	17.0 ^b	14.3 ^c	13.8 ^c	0.25
C18:0	9.42 ^a	8.71 ^a	6.61 ^b	3.00 ^c	3.73 ^c	0.15
C20:0	5.77 ^a	4.88 ^a	2.31 ^b	2.20 ^b	1.01 ^c	0.33
C22:0	0.95 ^a	0.44 ^b	0.37 ^b	0.30 ^b	0.29 ^b	0.02
C14:1c	1.44 ^c	3.08 ^a	3.71 ^a	3.88 ^a	3.90 ^a	0.12
C16:1c	2.01 ^b	2.21 ^b	2.93 ^b	3.18 ^a	3.51 ^a	0.03
C18:1c	10.4 ^c	19.0 ^b	19.2 ^b	25.8 ^a	26.1 ^a	0.74
C18:1n9t	1.00 ^c	1.72 ^b	1.88 ^b	1.93 ^b	2.11 ^a	0.02
C18:1n9c	0.82	0.86	0.80	0.83	0.87	0.17
C:22:1	0.10 ^b	0.67 ^a	0.71 ^a	0.76 ^a	0.98 ^a	0.02
C18:2n6	15.1 ^c	19.6 ^b	26.5 ^a	27.1 ^a	28.7 ^a	0.87
C20:5n3	0.97 ^c	1.60 ^b	1.72 ^b	1.88 ^b	2.77 ^a	0.10
C18:3n3	6.00 ^c	9.78 ^b	10.0 ^b	17.4 ^a	18.0 ^a	0.19
C20:4n6	3.11 ^b	3.40 ^b	3.59 ^b	3.80 ^b	7.22 ^a	0.22
C20:3n6	0.77 ^b	1.18 ^a	1.22 ^a	1.25 ^a	1.30 ^a	0.03
C22:6n3	0.98 ^c	2.05 ^b	2.10 ^b	2.18 ^b	3.18 ^a	0.42
TSFA ¹	46.14 ^a	35.95 ^b	30.64 ^b	24.00 ^c	22.65 ^c	0.03
TUFA ²	42.70 ^c	65.15 ^b	74.36 ^b	80.09 ^a	89.24 ^a	0.25

MUFA ³	15.77 ^c	27.54 ^b	29.23 ^b	30.08 ^b	35.07 ^a	0.44
PUFA ⁴	26.93 ^c	37.61 ^b	45.13 ^b	53.01 ^a	54.17 ^a	0.96
n-6:n-3 ⁵	1.38 ^b	1.80 ^b	2.41 ^a	2.89 ^a	3.65 ^a	0.03
Ant. Index ⁶	0.92 ^a	0.46 ^b	0.38 ^b	0.32 ^b	0.26 ^c	0.01

¹Total saturated fatty acid= C12:0 + C14:0 + C16:0 + C18:0 + C20:0 + C22:0

²Unsaturated fatty acid = (3 + 4)

³Mono unsaturated fatty acid= C14:1_c + C16:1_c + C18:1_c + C18:1n_{9t} + C18:1n_{9c} + C22:1

⁴Polyunsaturated fatty acid = C18:2 n₆ + C20:5 n₃ + C18:3n₃ + C20:4n₆ + C20:3n₆ + C: 22:6n₃

⁵n-6: n-3 = (C18:2 n₆ + C20:4n₆ + C20:3n₆) / (C20:5n₃ + C18:3n₃ + C: 22:6n₃), ⁶Antherogenic index = (C12:0+ 4×C14:0+ C16) / Σ of UFA

3. Results and Discussion

Table 1 reveals the nutrient content of the experimental diet, the chemical component contained crude protein (16.44%), crude fibre (8.82 %), ether extract (2.46 %), calcium (1.34 %), phosphorus (0.50 %) and metabolizable energy (2500.6 Kcal/kg). Higher crude fibre (12.08 %) was reported by Salisu and Kim [25] when carrot leaf meal was included in the diets of growing rabbits. According to Alinnor and Oze [26] adequate dietary intake of fibre can lower the risk of coronary heart disease and improve digestion in animals. Similarly Unigwe, *et al.* [27] reported a crude protein of crude protein range of 15.83 – 16.38 % in rabbits fed diets supplemented with dried neem leaf meal. However, all the values are within the nutritional requirement for rabbits as recommended by National Research Council [1].

The phytochemical components of *Albizia lebeck* seed in Table 2 reveals the presence of tannins, saponins, alkaloids, oxalate and glycosides. According to Hyun, *et al.* [5], a combination of multiple phytochemicals exerts synergistic effects to reduce negative effect of enteric infections. For instance, phenols have been reported to perform the role of antioxidants [28] tannins possess antimicrobial and antiviral activity [29]. Alkaloids and flavonoids have diuretic, antispasmodic, anti-inflammatory and analgesic effects [30]. According to Gadde, *et al.* [31], the main bioactive compounds of phytochemicals are polyphenols and their composition and concentration vary according to the plant, parts of the plant, harvesting season, geographical origin, environmental factors, storage conditions and processing techniques. However, the phytochemical values were below the lethal dose recommended by Alagbe and Grace [32]. This means that the feed is non-toxic and will not adversely affect the health of the animals; it further confirms the reports of Hawary, *et al.* [33] who stated that *Albizia lebeck* oil contains minerals, vitamins, essential and non-essential amino acids, which makes it useful in livestock feeds.

Table 3 shows fatty acid profile of rabbit meat supplemented with different levels of *Albizia lebeck* oil. Lauric acid (1.82 – 3.88 %), myristic acid (2.00 – 3.22 %), palmitic acid (13.8 – 22.9 %), stearic acid (3.73 – 9.42 %), arachidic acid (1.01 – 5.77 %), behenic acid (0.29 – 0.95 %), myristoleic acid (1.44 – 3.90 %), palmitoleic acid (2.01 – 3.51 %), linoleic acid (10.4 – 26.1 %), oleic acid (1.00 – 2.11 %), elaidic acid (0.80 – 0.87 %), linolelaidic acid (15.1 – 28.7 %), erucic acid (0.10 – 0.98 %), eicosapentenoic acid (0.97 – 2.77 %), α – linolenic acid (6.00 – 18.0 %), arachidonic acid (3.11 – 7.22 %), dihomoammalinolenic acid (0.77 – 1.30 %), docosahexenoic acid (0.98 – 3.18 %), total fatty acid ranges between (26.58 – 44.71 %), total unsaturated fatty acid (41.47 – 77.87 %), monounsaturated fatty acid (19.37 – 31.57 %) and polyunsaturated fatty acid (22.37 – 46.30 %) respectively. There were significant differences ($P < 0.05$) in the values obtained for TSFA, TUFA, MUFA and PUFA. TSFA value statistically reduced as the level of ALO increased in the diet. This confirms the earlier report by Alagbe and Grace [32] that ALO has the ability to reduce the cholesterol level in broiler chicks; this could be attributed to the presence of phytochemicals contained in ALO seeds. Hawary, *et al.* [33], also reported that ALO is rich in protein, essential and non – essential acid as well as PUFA. This is a clear indication that ALO has the ability to reduce SFA which in turn will improve the nutritive value of meat and reduce the incidence of cardiovascular infection. The result is consistent with the findings of Kamran, *et al.* [16] who reported that SFA level in the meat of rabbits decreased by feeding diets enriched with flaxseed oil. PUFA and n-6: n-3 level increased as the level of ALO increased especially at 0.4 %, thus ALO has the ability to modulate the fatty acid of meat. According to Katalin and Loana [34] omega – 3 and omega -6 polyunsaturated fatty acids performs multiple biological roles such as influencing the inflammatory cascade, reducing oxidative stress, presenting neuro- protection and cardiovascular protection. This result is in agreement with the findings of Suriya, *et al.* [35] and Trebusak, *et al.* [36] when linseed was supplemented in the diets of rabbits. However, omega -3 and omega - 6 polyunsaturated fatty acid ratios was within the range recommended by Simopoulos [37]. Antherogenic index is a parameter used to determine meat safety, the lower the value the safer the meat. Supplementation of ALO at 0.4 % significantly reduced the antherogenic index when compared with animals fed 0% ALO. This reveals that ALO can reduce the incidence of arteriosclerosis and other cardiovascular infections which is in accordance with the reports of Kholif, *et al.* [38].

4. Conclusion

It can be concluded that ALO has proved their efficacy to perform multiple biological activities due to the presence of bioactive chemicals or secondary metabolites. Supplementation of ALO at 0.4 % in the diet of rabbits modulates the fatty acid composition and nutritive value of meat. High PUFA composition in rabbit meat prevents cardiovascular and other degenerative diseases therefore bridging the gap between food safety and livestock production.

References

- [1] National Research Council, 1977. *Micro livestock: Little known small animals with promising economic future XVII*. Washington D.C.: National Academy Press. p. 449.
- [2] Oyen, L. P. A. and Dung, N. X., 1999. *Plant resources of South East Asia no. 19. Essential oil plants*. Backhuys Publishers.
- [3] Enzo, P. N., 2007. "Comparative antioxidant capacity, membrane stabilization, polyphenol composition and cytotoxicity of the leaf and stem of *Cissu multistriata*." *African Journal of Biotechnology*, vol. 7, pp. 3129-3133.
- [4] Mohammed, F., Singh, P. P., and Irchhaiya, R., 2012. "Review on Albazia lebbeck a potent herbal drugs." *International Research Journal of Pharmacy*, vol. 3, pp. 11-15.
- [5] Hyun, L., Yanhong, L., Sergio, C., Mariano, E. O., Fang, C., Ron, L. C., Sungtaek, O., and Cyril, G. G., 2018. "Phytochemical as antibiotic alternatives to promote growth and enhance host health." *Journal of Biomedical Sciences*, vol. 10, pp. 49-76.
- [6] William, P., Stirling, E., and Keynes, N., 2004. "Food fears: a natural survey on the attitudes of Australian adults about the safety and quality of food." *Asian Pacific Journal of Clinical Nutrition*, vol. 13, pp. 32-39.
- [7] Lee, K., Everts, H., and Beynen, A., 2004. "Essential oil in broiler nutrition." *International Journal of Poultry Science*, vol. 3, pp. 738-752.
- [8] Gopal, R. S. and Asmita, M. P., 2014. "Effect of Saponin of *Albizia lebbeck* Benth on the reproductive performance of male albino rats." *Journal of Ethnopharmacology*, vol. 96, pp. 31-36.
- [9] Everist, F. L., 1986. "Antioxidant activity and phenolic compounds in 32 selected herbs." *Food Chem.*, vol. 105, pp. 940-949.
- [10] Waheed, S., Khalid, B., and Ashraf, C. M., 2000. "Chemical analysis of *Albizia lebbeck*." *Journal of Chemistry Society. Pak.*, vol. 22, p. 56.
- [11] Kirsti, T., Kettunen, H., Bento, M. H., Saarienen, M., Lahtinen, S., and Routonen, N., 2010. "The effect of feeding essential oil on broiler performance and gut microbiota." *British Poultry Science*, vol. 51, pp. 381-392.
- [12] Bourre, J., 2004. "Effect of increasing the omega -3 fatty acid in the diets of animals on the animal products consumed by humans." *Journal of Medicinal Sciences*, vol. 21, pp. 773-779.
- [13] Alagbe, J. O., Shittu, M. D., and Eunice, A. O., 2020. "Prospect of leaf extracts on the performance and blood profile of monogastric – A review." *International Journal of Integrated Education*, vol. 3, pp. 122-127.
- [14] Alagbe, J. O., 2019. "Effects of dried *Centella asiatica* leaf meal as a herbal feed additive on the growth and serum biochemistry of broiler chicken." *International Journal of Animal Research*, vol. 3, pp. 1-12.
- [15] Trebusak, V. U., Chukwu, G. C., and Adeolu, A. I., 2011. "Effect of ground ginger and garlic on the growth Performance, carcass quality and economics of production of broiler chickens." *Global Journal of Bio-Science and Biotechnology*, vol. 3, pp. 225-229.
- [16] Kamran, K., Sarzamin, K., Nazir, A. K., and Naseer, A., 2018. "Effect of flaxseed supplementation of feed on growth, carcass yield, meat and fatty acid profile of rabbit carcass." *Pakistan Journal of Zoology*, vol. 50, pp. 2037-2043.
- [17] Busquet, M., Calsamiglia, S., Ferret, A., Carro, M. D., and Kamel, C., 2005. "Effect of garlic oil and four of its compound on rumen microbial fermentation." *Journal of Dairy Science*, vol. 88, pp. 4393-4404.
- [18] Cerisuelo, A., Marin, C., Sanchez, V. F., Gomez, E. A., and Duran, F., 2014. "The impact of a specific blend of essential oil components and sodium butyrate in feed on the performance and *Salmonella* counts in experimentally challenged broilers." *Journal of Poultry Science*, vol. 93, pp. 599-606.
- [19] Folch, J., Lees, M., and Sloane-Stanley, G., 1957. "A simple method for the isolation and purification of total lipids from animal tissues." *Journal of Biological Sciences*, vol. 226, pp. 497-509.
- [20] Elshater, A. E., Muhammad, M. A. S., and Mahrous, M. A. M., 2009. "Effect of ginger extracts consumption on levels of blood glucose, lipid profile and kidney functions in alloxane induced-diabetic rats." *Egypt. Acad. J. Biol. Sci.*, vol. 2, pp. 153-162.
- [21] Sofowora, A., 1993. *Medicinal plants and Traditional Medicine in Africa*. Nigeria: Spectrum Books Limited Ibadan. p. 289.
- [22] Harbone, I. B., 1973. *A guide to modern techniques to plant analysis*. 2nd ed. New York, USA: Chapman and hall.
- [23] Boham, B. A. and Kocipai, A. C., 1974. "Flavonoids and condensed tannins from the leaves of Hawaiian *vaccinium vaticulatum* and *V. calycinium*." *Journal of Poultry Science*, vol. 48, pp. 458-463.
- [24] Duncan, D. B., 1955. "Multiple range and multiple F-test." *Biometrics*, vol. 11, pp. 1-42.
- [25] Salisu, D. U. and Kim, I. H., 2012. "Efficacy of phytogetic feed additive on performance production and health status of monogastric animals." *A review. Annals of Animal Science*, vol. 17, pp. 929-948.
- [26] Alinnor, J. and Oze, R., 2011. "Chemical Evaluation of the Nutritive value of *Pentaclethra macrophylla* benth (*African Oil Bean*) Seeds." *Pakistan Journal of Nutrition*, vol. 10, pp. 355-359.
- [27] Unigwe, C. R., Balogun, F. A., Okarafor, U. P., Idah, I. S., Abonyi, F. O., and Olona, J. F., 2016. "Effect of neem leaf (*Azadirachta indica*) meal on growth performance abd haematology of rabbits." *World Scientific News*, vol. 55, pp. 51-62.
- [28] Andjelkovic, P., Charu, G., and Girish, S., 2016. "Importance of phytochemicals in nutraceuticals." *Journal of Chinese Medicine Research and Development*, vol. 1, pp. 70-78.

- [29] Adisa, R. M., Choudhary, E. A., Adenoye, G. A., and Olorunsogo, O. O., 2010. "Hypoglycaemic and biochemical properties of *Cnestis ferruginea*." *African Journal of Traditional Complementary Alternative Medicine*, vol. 7, pp. 185-194.
- [30] Ujowundu, C. O., Okafor, O. E., Agha, N. C., Nwaogu, L. A., Igwe, K. O., and Igwe, U. C., 2010. "Phytochemical and chemical composition of *Combretum zenkeri* leaves." *Journal of Medicinal Plants Research*, vol. 4, pp. 965-96.
- [31] Gadde, M. F., Khairunnesa, M., and Das, S. C., 2017. "Use of non-antibiotic growth promoter "Grow Power" in commercial broiler diet." *Bangladesh Journal of Animal Science*, vol. 44, pp. 33-39.
- [32] Alagbe, J. O. and Grace, F. R., 2019. "Effect of *Albizia* seed oil dietary supplementation on the haematological and serum biochemical parameters of weaner rabbits." *Sumerianz Journal of Agriculture and Veterinary*, vol. 2, pp. 96-100.
- [33] Hawary, S., El-Fouly, K., Sokkar, N. M., and Talaat, Z., 2011. "A phytochemical profile of *albizia lebbek* (L.) benth. Cultivated in Egypt." *Asian Journal of Biochemistry*, vol. 6, pp. 122-141.
- [34] Katalin and Loana, 2017. "Nutritional evaluation of chemical component of leaves stalks and stems of sweet potatoes (*Ipomea batatas*)." *Food Chemistry*, vol. 68, pp. 359-367.
- [35] Suriya, K. R., Idrus, Z., Nordiana, A. A., Mahdi, E., and Goy, Y. M., 2014. "Effects of two herbal extracts and Virginiamycin supplementation on the growth performance, intestinal microbial population and fatty acid composition of broiler chickens." *Asian Australian Journal of Animal Science*, vol. 27, pp. 375-382.
- [36] Trebusak, T., Leva, A., Frankic, T., Salobir, J., and Pirman, T., 2014. "Effect of dietary linseed oil and *Ganoderma lucidum* or olive leaves supplementation on fatty acid composition and oxidative status of rabbits." *World Rabbit Science*, vol. 22, pp. 71-81.
- [37] Simopoulos, A. P., 2001. "N-3 fatty acids and human health: defining strategies for public policy." *Journal of Animal Science*, vol. 36, pp. 83-89.
- [38] Kholif, A. E., Gouda, G. A., Olafadehan, O. A., and Abdo, M. M., 2017. "Effect of replacement of *Moringa oliferi* for berseem clover in the diets of Nubian goats on feed utilisation, milk yield composition and fatty acid profile." *The Animal Consortium Journal*, vol. 12, pp. 964-972.