



An Analysis of the Impact of Various Sampling Designs on the Headcount Index: A Simulation Study Based on the EU-SILC

Pablo J. Moya Fernández (Corresponding Author)

Department of Applied Economics, Faculty of Social and Legal Sciences of Melilla, University of Granada, Spain

Email: pimoyafernandez@ugr.es

Encarnación Álvarez

Department of Quantitative Methods in Economics and Business, Faculty of Economics and Business, University of Granada, Spain

Ángela González Medina

Department of Quantitative Methods in Economics and Business, Faculty of Economics and Business, University of Granada, Spain

Article History

Received: September 20, 2020

Revised: October 19, 2020

Accepted: October 26, 2020

Published: October 31, 2020

Copyright © 2020 ARPG &
Author

This work is licensed under the
Creative Commons Attribution
International



CC BY: Creative
Commons Attribution License
4.0

Abstract

The analysis and the comparison of poverty between regions and countries are important topics in social sciences, which have relevant demands of many national (Cáritas, Intermón Oxfam, Cruz Roja, etc) and international (UN, World Bank, OECD, Eurostat, IMF, etc) agencies and organizations. One of the most common poverty indicators in practice is the headcount index, which analyzes the proportion of individuals considered as poor in a population. In this paper, we first analyze the impact on the headcount index when different sampling designs are considered. Note that this study is based on real data sets taken from different countries of the European Union, and the empirical measures for comparisons are based on different Monte Carlo simulation studies. For instance, we observe that stratified sampling has the best performance in comparison to alternative sampling designs. Post-stratification performs similar to simple random sampling without replacement, and the use of auxiliary information provides similar results to ones derived from stratified sampling. Second, we also analyze the empirical performance of different variance estimators under the commented sampling designs. We conclude that they have a similar empirical performance, and they provide, in general, confidence intervals with desirable coverage rates.

Keywords: Poverty line; Proportion; Random sample; Monte Carlo simulation; Confidence interval.

1. Introduction

The main aim of this manuscript is to analyze the impact of various common sampling designs on the estimation of the proportion of individuals classified as poor. This proportion is also named as the headcount index or low income proportion. Note that this study is based on various real data sets extracted from the 2011 European Union Survey on Income and Living Conditions (EU-SILC). The EU-SILC is carried out by the statistical agency Eurostat. Some references that discuss about the EU-SILC are Eurostat (2010), Eurostat (2013), Muñoz *et al.* (2015), López-Escobar *et al.* (2016), Berger and Torres (2014), Goedemé (2010), Goedemé (2013) and Van Kerm (2007), Museux (2005), among others.

One of the main challenges from governments and national and international R&D+i strategies is to eradicate poverty. In addition, many international (UN, World Bank, OECD, Eurostat, IMF, etc) and national (Cáritas, Intermón Oxfam o Cruz Roja, etc) organizations require a better understanding of the poverty. In this sense, first of all, this paper is motivated by the fact that headcount index is an important topic with an increased interest from all these institutions. Also statistical agencies show high interest in this field of study. For example, Eurostat (2010), Eurostat (2013), and Haughton and Khandker (2005) obtain regularly estimations of the headcount index.

In order to study the different situations that derivate into a poverty position, it is necessary to make analysis from information and indicators that allow an accurate and reliable valuation from the reality. For this reason, it is crucial to have the proper instruments to make a right approximation from this problem and therefore the right decision making in the political sphere. Although plentiful literature does exist about the poverty analysis and the use of indicators, nowadays there are no studies that focus on the impact of different sampling designs over one of the most important poverty indicators, the headcount index.

Therefore, this study is based in the intense debate about the necessity of introducing improvements in the measurement, estimation and analysis stage (Eurostat, 2013; Goedemé, 2013). This problem has its root in the fact that the majority of the social and economic indicators have to be formed from sampling information (see Haughton and Khandker (2005); Eurostat (2010)), which means, there is a necessity of using estimation methods that generate more accurate and high quality estimations.

It aims to provide a reference for the measurement and the estimation of the headcount index through different estimations methods and a basis for the improvement of the national poverty reduction strategies and policies as well.

We consider different sampling designs commonly used in practice, such as simple random sampling without replacement (SRSWOR), simple random sampling and using auxiliary information, stratified random sampling, and post-stratified sampling. The point estimation of the headcount index is an important topic, but sampling errors may arise in practice and it is quite important to measure the level of this sampling error, and this is the purpose of the variance estimation of a given estimator. For this reason, we also calculate the variance estimation of the different estimators based on the corresponding sampling designs. Finally, note that Monte Carlo simulation studies are used to analyze the precision of the different estimators and variance estimators. The analysis of the impact of the various sampling designs can be also realized. In particular, estimators are compared in terms of Relative Bias (RB) and Relative Root Mean Square Error (RRMSE). The precision of the variance estimators is analyzed via the RB and the coverage of the corresponding confidence intervals. For instance, such empirical measures are defined and described by Chambers and Dunstan (1986), Rao *et al.* (1990), Silva and Skinner (1995), Harms and Duchesne (2006), Berger and Munoz (2015), Muñoz *et al.* (2015), López-Escobar *et al.* (2016), etc.

2. The Headcount Index

From a social point of view, the poverty is a complex phenomenon, in which have influenced many factors and that should appear on the political and economic agenda. Ravallion (1999) “A credible measure of poverty can be a powerful instrument for focusing the attention of policy makers on the living conditions of the poor; it is easy to ignore the poor if they are statistically invisible.”

Some of the authors that specialize in this topic are, among others, Sen (1976), Ravallion (1999), Foster (1998), Foster *et al.* (1984), Khandker (2005), Haughton and Khandker (2009), Atkinson (1987).

Others important reasons to measure the poverty, in spite of its complexity, are the following. It is necessary to know the characteristics and behaviors of the populations in which the poverty correcting measures should be implemented. By getting this aim, the effect or impact of the measures could be greater. Also predicting the effects of poverty projects and policies with the aim of improving or cancel them if they are not viable. Finally, we can point out that evaluating the actions from governments and institutions is not possible without measures that summarize in a value the information provides by data.

To sum up, the poverty measurement as well as the analysis and evaluation of programs and projects are decisive to set policies to reduce the poverty worldwide. In this sense, the use of accurate indicators is important in order to explain the distribution of poor people and the real proportion from these.

Poverty as a pronounced deprivation in wellbeing (Haughton and Khandker, 2005) can be studied from many different perspectives. So, there are many ways to measure it, starting from the different definitions. In this paper, we focus in the monetary approach, which is the most use to compute poverty and attributes a monetary value to poverty. For this, it is necessary to establish a poverty line or income threshold. Those individuals with income below the line are deemed to be poor. From this point of view, the measurement of poverty is, therefore, a kind of measurement of the income necessary to meet certain basic needs (poverty line), including food and nonfood needs (Haughton and Khandker, 2009).

The relative poverty line is defined as a point in the distribution of income or expenditure and, hence, the line can be updated automatically over time for changes in living standards (Zheng, 2001). In practice, researchers often specify the relative poverty line as a percentage of mean or median income or expenditure. For instance, EUROSTAT uses a relative poverty line given by the 60 % of the national median equivalised disposable income. This means that the threshold below which a person is qualified as poor, is set at 60 % of the national median equivalised disposable income. The OECD used as poverty line the 50% of the national median equivalised disposable income. Regarding relative poverty lines see also, Thompson (2013).

After defining a poverty line, we need a statistical measure that assigns a value to the distribution of poor. There exist a wider set of poverty measures (Haughton and Khandker, 2005). The measure used in this paper is the headcount index, which gives us the proportion of people below an official poverty line or, the proportion of people classified as poor into a population (Álvarez *et al.*, 2014). This parameter is given by P , because we are talking about a proportion, so formally we obtain that,

$$P = N_p/N$$

is the headcount index for a given population, where N_p , is the number of poor into this population, and N is the population size.

The greatest virtues of the headcount index are that it is simple to construct and easy to understand; and these are important qualities (Haughton and Khandker, 2009). Such authors also point out that this index has some limitations as i) does not take the intensity of poverty into account; ii) does not indicate how poor the poor are, and hence does not change if people below the poverty line become poorer; or iii) the estimates should be calculated for individuals, not households.

3. Some Sampling Designs and Estimators of the Headcount Index

The purpose of this paper is to analyze the estimation of the headcount index under different sampling designs. In particular, we can compare sampling designs based on stratification to sampling designs without stratification. In addition, we can analyze numerically the gain of using auxiliary information at the estimation stage with real data

taken from different countries from the European Union. The idea is to observe numerically the effect on this parameter when different estimation methods are applied. For this reason, in this section we describe various known estimation methods which could be used in practice.

In the previous section, we commented that the headcount index for a given population is denoted as P . This parameter is unknown since it is not possible to know the value of the variable of interest y (the equivalised net income) at the population level. To remedy this, we can select a sample (which is denoted as s), with size n , from this population, and where the variable of interest can be observed for the individuals selected in this sample. $\{y_1, \dots, y_n\}$ denote the sampled values of the variable of interest. The various sampling designs considered in this study are briefly described as follows.

3.1. Simple Random Sampling without Replacement

The simple random sampling without replacement (Cochran, 1977; Särndal *et al.*, 2003), SRSWOR, has the property that all the individuals in the population have the same probability of selection. In this situation, we get a sample or subset of individuals, where each one is chosen randomly and each individual has the same probability of being chosen at any stage during the sampling process, being this statement its main principle. This method is developed here without replacement, which avoids choosing any individual more than once. Note that SRSWOR is the sampling method most commonly used, and it is considered as basic type of sampling, since it can be a component of other more complex sampling methods.

Assuming that the poverty line is given by L , the estimator of the headcount index under SRSWOR can be expressed as:

$$\hat{p} = \frac{n_p}{n} = \frac{1}{n} \sum_{i \in s} \delta(y_i \leq L),$$

where

$$n_p = \sum_{i \in s} \delta(y_i \leq L)$$

is the number of poor in the sample s , and $\delta(\cdot)$ is the indicator variable, which takes the value 1 if its argument is true and takes the value 0 otherwise. It is well known that \hat{p} is an unbiased estimator, i.e., $E(\hat{p}) = P$. The problem of estimating the variance of estimators is discussed in Section 3.5.

3.2. Simple Random Sampling with Auxiliary Information

In sample surveys, auxiliary information (Berger and Munoz, 2015; Chambers and Dunstan, 1986; Dorfman, 2009; Muñoz *et al.*, 2015; Rao *et al.*, 1990; Rao *et al.*, 1992; Rueda and Muñoz, 2011; Silva and Skinner, 1995; Wu, 1999) is regularly used to increase the precision of estimators, and most commonly when the problem is the estimation of means or totals (Breidt and Opsomer, 2000; Chen and Sitter, 1999; Deville and Särndal, 1992).

There exist many methods based on auxiliary information, but some of the most known are the ratio and regression techniques. For simplicity, we consider the ratio method. Let x be an auxiliary variable related to the variable of interest y . Taking into account our data base EU-SILC, which provides information on income, poverty, social inclusion and living conditions for a sample of households and individuals; in our context, the auxiliary variable can be the wages or wage inequality, tax on income contributions and so on (Berger and Munoz, 2015; Muñoz *et al.*, 2015; Rueda and Muñoz, 2011).

The ratio estimator of the headcount index can be calculated as

$$\hat{p}_r = \frac{\hat{p}}{\hat{p}_x} P_x$$

where

$$\hat{p}_x = \frac{1}{n} \sum_{i \in s} \delta(\hat{R}x_i \leq L)$$

is the customary estimator of

$$P_x = \frac{1}{N} \sum_{i \in U} \delta(\hat{R}x_i \leq L)$$

and

$$\hat{R} = \frac{\bar{y}}{\bar{x}}$$

is the customary estimator of the population ratio

$$R = \frac{\bar{Y}}{\bar{X}}$$

where \bar{y} and \bar{x} are, respectively, the sample means of the interest and auxiliary variables, and \bar{Y} and \bar{X} are the corresponding population means. It is well known that the ratio estimator is a biased estimator, but it is asymptotically unbiased.

3.3. Stratified Simple Random Sampling

The stratified sampling (Cochran, 1977; Särndal *et al.*, 2003), is a very popular sampling method in which the first step is to divide the whole population into different groups. Hence, the population of N units is first divided into H subpopulations called “strata”. Then, we consider that the sample is selected randomly through SRSWOR from

each stratum, so this whole procedure is named as stratified simple random sampling (SSRS). Strata can be formed by using many methods, but it is most common and simple to use a variable such as age, gender, socioeconomic status, geographic region, religion, nationality, educational attainment, etc. Note that the customary way to carry out this method is to estimate the parameters separately for each stratum, and then to combine them into a weighted estimator. The estimator of the headcount index based upon SSRS is given by

$$\hat{p}_{SSRS} = \frac{1}{N} \sum_{h=1}^H N_h \hat{p}_h \quad (1)$$

where N_h is the population size of the h th stratum, and

$$\hat{p}_h = \frac{1}{n_h} \sum_{i=1}^{n_h} \delta(y_{hi} \leq L),$$

is the estimation of the headcount index for the h th stratum, and n_h is the sample size for the h th stratum.

In this paper we consider the method “4E” suggested by [Silva and Skinner \(1995\)](#) to obtain the strata, and which provided good results to such authors. This method consists on ordering the auxiliary variable and take strata with equal (E) number of units according to this auxiliary variable. We consider a total of 4 strata, hence this method is named as 4E. The sample sizes are obtained by using proportional allocation (see also [Silva and Skinner \(1995\)](#)).

3.4. Post-Stratified Sampling

Post-stratification sampling consists on applying the concept of stratification according to auxiliary information about the sampled population but after the selection of the sample. Some relevant references that discuss about post-stratification are [Särndal et al. \(2003\)](#), [Silva and Skinner \(1995\)](#) and [Valliant \(1993\)](#), among others.

The expression of the estimator of the headcount index under post-stratified sampling is the same than the expression (1) given by SSRS. This estimator will be denoted as $\hat{p}_{P,ST}$. However, the difference is the fact that the stratification is carried out after the sample is selected. It can be interesting to observe the differences in accuracy when the stratification is considered before and after the sample is selected. This study can be also seen in Section 4. Note that post-strata are obtained by using the method 4E described in Section 3.3 and suggested by [Silva and Skinner \(1995\)](#).

3.5. Variance Estimator

We have previously defined point estimators for the headcount index. However, the problem of estimating the variance of the estimator of the headcount index is also an important topic, since we can have more information about the accuracy of the given estimation. An additional aim in this study is to analyze and compare various variance estimators under the sampling designs under study. In particular, we analyze the Horvitz-Thompson variance estimator, the Sen-Yates-Grundy variance estimator ([Sen, 1953; Yates and Grundy, 1953](#)), and the Hájek variance estimator ([Hájek, 1964](#)) for the Narain-Horvitz-Thompson ([Horvitz and Thompson, 1952; Narain, 1951](#)) point estimator for the headcount index. Such expressions can be also seen in [López-Escobar et al. \(2016\)](#), and which create an R package to calculate such expressions in practice.

3.6. The Horvitz-Thompson Variance Estimator

For a general sampling design, the Horvitz-Thompson variance estimator of the estimator of a population mean is given by

$$\hat{V}_{HT}(\bar{y}) = \frac{1}{N^2} \sum_{i \in S} \sum_{j \in S} \frac{\pi_{ij} - \pi_i \pi_j}{\pi_{ij}} \frac{y_i}{\pi_i} \frac{y_j}{\pi_j} \quad (2)$$

where π_i and π_{ij} are, respectively, the first and the second order inclusion probabilities. Assuming SRSWOR these quantities are given by

$$\pi_i = \frac{n}{N}; \quad \pi_{ij} = \frac{n(n-1)}{N(N-1)}.$$

For the case of estimating the headcount index, the corresponding Horvitz-Thompson variance estimator is obtained by substituting y_i by $\delta(y_i \leq L)$ into equation (2), i.e.,

$$\hat{V}_{HT}(\hat{p}) = \frac{1}{N^2} \sum_{i \in S} \sum_{j \in S} \frac{\pi_{ij} - \pi_i \pi_j}{\pi_{ij}} \frac{\delta(y_i \leq L)}{\pi_i} \frac{\delta(y_j \leq L)}{\pi_j}. \quad (3)$$

Similarly, we can obtain the Horvitz-Thompson variance estimator of the ratio estimator of the headcount index. In this situation, the corresponding expression is given by

$$\hat{V}_{HT}(\hat{p}_r) = \frac{1}{N^2} \sum_{i \in S} \sum_{j \in S} \frac{\pi_{ij} - \pi_i \pi_j}{\pi_{ij}} \frac{z_i}{\pi_i} \frac{z_j}{\pi_j},$$

where

$$z_i = \delta(y_i \leq L) - \frac{\hat{p}}{\hat{p}_x} \delta(\hat{R}x_i \leq L)$$

Assuming SSRS, the Horvitz-Thompson variance estimator of the estimator of the headcount index is given by

$$\hat{V}_{HT}(\hat{p}_{SSRS}) = \frac{1}{N^2} \sum_{h=1}^H N_h^2 \hat{V}(\hat{p}_h)$$

where $\hat{V}(\hat{p}_h)$ is the variance given by the expression (3) but calculated at the h th stratum.

Finally, the Horvitz-Thompson variance estimator of the estimator of the headcount index under post-stratification is given by

$$\hat{V}_{HT}(\hat{p}_{P.ST}) = \frac{1}{N^2} \sum_{i \in s} \sum_{j \in s} \frac{\pi_{ij} - \pi_i \pi_j}{\pi_{ij}} \frac{u_i}{\pi_i} \frac{u_j}{\pi_j},$$

where

$$u_i = \delta(y_i \leq L) - \hat{p}_h$$

3.7. The Sen-Yates-Grundy Variance Estimator

The Sen-Yates-Grundy variance estimator can be similarly obtained under the difference sampling designs. However, we have to substitute the expression (2) by

$$\hat{V}_{SYG}(\bar{y}) = \frac{1}{N^2} \frac{1}{2} \sum_{i \neq j} \frac{\pi_i \pi_j - \pi_{ij}}{\pi_{ij}} \left(\frac{y_i}{\pi_i} - \frac{y_j}{\pi_j} \right)^2 = \sum_{i < j} \frac{\pi_i \pi_j - \pi_{ij}}{\pi_{ij}} \left(\frac{y_i}{\pi_i} - \frac{y_j}{\pi_j} \right)^2.$$

For example, the corresponding Sen-Yates-Grundy variance estimator of the headcount index under SRSOWR is given by

$$\hat{V}_{SYG}(\hat{p}) = \frac{1}{N^2} \frac{1}{2} \sum_{i \neq j} \frac{\pi_i \pi_j - \pi_{ij}}{\pi_{ij}} \left(\frac{\delta(y_i \leq L)}{\pi_i} - \frac{\delta(y_j \leq L)}{\pi_j} \right)^2.$$

Expressions for alternative estimation methods are similarly defined.

3.8. The Hájek Variance Estimator

Finally, we also compute in this paper the Hájek variance estimator (Hájek, 1964). For the problem of estimating a population mean, this variance estimator is given by

$$\hat{V}_H(\bar{y}) = \frac{n}{N^2(n-1)} \left[\sum_{i \in s} \left(\frac{y_i}{\pi_i} \right)^2 (1 - \pi_i) - \hat{d} \hat{G}^2 \right],$$

where

$$\hat{d} = \sum_{i \in s} (1 - \pi_i)$$

and

$$\hat{G} = \frac{1}{\hat{d}} \sum_{i \in s} \frac{(1 - \pi_i) y_i}{\pi_i}.$$

Following the notation used for the Horvitz-Thompson variance estimator, the Hájek variance estimator of estimators of the headcount index can be easily defined. For this purpose, we just need to replace y_i by $\delta(y_i \leq L)$ into the previous expressions.

4. Monte Carlo Simulation Studies

In this section, we carry out various Monte Carlo simulation studies which analyze the aspects described in this paper. Note that the various studies are based on real data sets extracted from the 2011 EU-SILC, and which are taken from 8 countries from the European Union. In particular, we analyze numerically the bias and the efficiency of estimators of the headcount index when samples are selected under different sampling designs. In addition, we evaluate different variance expressions under the same sampling designs. For this purpose, the empirical coverages of the corresponding confidence intervals are computed.

4.1. Populations

The simulation studies are based on real survey data obtained from the 2011 EU-SILC. We considered data from 8 different countries and they were considered because they have different properties according to various poverty indicators, i.e., they have quite different values for the headcount index, the poverty gap index, etc. Such countries considered in this paper are: Belgium, Bulgaria, Spain, Slovenia, Italy, Lithuania, Poland and UK. Note that the variable of interest, y , is equivalised net income, whereas we consider the tax on income contributions as the auxiliary variable. Note that this choice is due to the fact that both variables have a large linear relationship for the various populations used in this paper. We consider that the data associated to each country are considered as a population with size N , from which samples will be selected, and our studies can be thus realized.

4.2. Empirical Measures

Monte Carlo simulation studies are based on different empirical measures, which are described in this section. Point estimators of the headcount index are compared in terms of Relative Bias (RB) and Relative Root Mean Square Error (RRMSE), which are defined, respectively, by:

$$RB[\tilde{p}] = \frac{E[\tilde{p}] - P}{P}$$

$$RRMSE[\tilde{p}] = \frac{\sqrt{MSE[\tilde{p}]}}{P},$$

where \tilde{p} denotes a given estimators of the parameter P , and $E[\cdot]$ and $MSE[\cdot]$ are, respectively, the empirical expectation and mean square error based on $R = 10000$ simulation runs, i.e.,

$$E[\tilde{p}] = \frac{1}{R} \sum_{r=1}^R \tilde{p}(r)$$

$$MSE[\tilde{p}] = \frac{1}{R} \sum_{r=1}^R (\tilde{p}(r) - P)^2$$

where $\tilde{p}(r)$ denotes the value of \tilde{p} at the r th simulation run.

On the other hand, the accuracy of the different variance estimators is analyzed via the corresponding relative bias of such variance estimators, and also, the coverage of confidence intervals. For the r th iteration run, the lower and the upper limits for the confidence interval of a given estimator \tilde{p} are defined, respectively, by:

$$L(r) = \tilde{p}(r) - 1.96 \sqrt{\hat{V}(\tilde{p}(r))}$$

$$U(r) = \tilde{p}(r) + 1.96 \sqrt{\hat{V}(\tilde{p}(r))}.$$

We considered confidence intervals with a 95% for the confidence level, and for this reason, the empirical coverages should take values close to the nominal level of 95%. The coverage rates at the 95% confidence level are defined as,

$$CR[\tilde{p}] = \frac{1}{R} \sum_{r=1}^R \delta(L(r) \leq P \leq U(r))$$

In addition, we computed the percentage of times that the real parameter is, respectively, below and above the lower and upper limits of the confidence intervals. Note that the ideal situation is to have percentages close to 2.5%.

4.3. Results - Empirical Comparison of Estimators

We now present results derived from the point estimation of the headcount index. The relative biases and efficiency of the different estimation methods are presented in Table 1. We can observe that the various estimators have a good performance in terms of RB, since such values are smaller than 1% in absolute terms. We observe that the more efficient results are obtained under stratified simple random sampling, since the smallest values of RRMSE are obtained under this situation. Then, we observe that the ratio method also provides estimators with small values of RRMSE. Note that it is expected to improve such values in the case of population with larger values for the population correlation coefficient. SRSWOR and post-stratification are the sampling designs with the worst performance in terms of efficiency. We expected better results when the estimator is based on post-stratification, but the linear correlation coefficient is no large, and this could be the reason of this poor performance.

Table-1. Values of RB (%) and RRMSE (%) for the different estimators of the headcount index P and using a sampling fraction of $f=1\%$ ($=n/N$). ρ is the population linear correlation coefficient between the auxiliary and interest variables

| Country | P | \hat{P}_{SRSWOR} | | \hat{P}_r | | \hat{P}_{SSRS} | | $\hat{P}_{P,ST}$ | |
|-----------|------|--------------------|-------|-------------|-------|------------------|-------|------------------|-------|
| | | RB | RRMSE | RB | RRMSE | RB | RRMSE | RB | RRMSE |
| Belgium | 0,65 | 0,0 | 24,9 | 0,0 | 23,6 | -0,1 | 22,5 | 1,0 | 25,1 |
| Bulgaria | 0,70 | 0,1 | 16,8 | 0,2 | 15,2 | 0,0 | 14,8 | 0,1 | 16,8 |
| Spain | 0,65 | -0,1 | 11,7 | 0,1 | 11,2 | -0,1 | 11,0 | 0,6 | 11,7 |
| Slovenia | 0,83 | 0,0 | 20,1 | -0,2 | 18,1 | -0,1 | 17,7 | 0,0 | 20,1 |
| Italy | 0,88 | -0,3 | 12,3 | -0,2 | 10,8 | 0,0 | 10,3 | -0,6 | 12,3 |
| Lithuania | 0,66 | -0,2 | 22,6 | 0,0 | 21,3 | 0,0 | 20,5 | -0,2 | 22,6 |
| Poland | 0,82 | 0,1 | 14,4 | 0,3 | 13,6 | -0,1 | 13,1 | 0,5 | 14,5 |
| UK | 0,83 | -0,1 | 21,4 | -0,1 | 19,5 | 0,2 | 19,1 | 0,9 | 21,6 |

The results related to the variance estimation of estimators of the headcount index are presented in Table 2-5. In particular, Table 2 presents the relative bias associated to such variance estimators. For a given sampling design, we first observe that the various variance estimation methods provide similar results. We observe that the post-stratification method has a large negative bias, with values larger than 30% in relative terms. For alternative sampling designs and for the different data sets, the relative biases associated to the variance estimators give reasonable results. Another important issue is the precision of confidence intervals, which is analyzed via the coverage rates (see Table 3). We observe how the bias found under the post-stratification sampling design gives confidence intervals with a poor performance, i.e, they have coverage rates about 90%, which is far away from the required nominal level of 95%. Alternative sampling designs give desirable coverage rates, close to the required 95%, and the different methods also give similar results. Finally, Tables 4 and 5 give the percentages of times that the real parameter is, respectively, below and above the lower and upper limits of confidence intervals. Note that it is desirable to obtain balanced values, but this is not the situation for the different examples. In general, we observe that values of Table 5 are slightly larger than values of Table 4, and this indicates that confidence intervals should have larger upper limits. In addition, Table 4 can have smaller values due to the fact that the parameter is a proportion close to 0, and the lower limits could be values very close to 0, hence it is quite difficult to find situations where the real headcount index is below such lower limits.

Table-2. Values of RB (%) for different variance estimators of the headcount index P and using a sampling fraction of $f=1\%$ ($=n/N$)

| Country | \hat{P}_{SRSWOR} | | | \hat{P}_r | | | \hat{P}_{SSRS} | | | $\hat{P}_{P.ST}$ | | |
|-----------|--------------------|-----------------|-------------|----------------|-----------------|-------------|------------------|-----------------|-------------|------------------|-----------------|-------------|
| | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H |
| Belgium | 0,5 | 0,5 | 0,5 | -3,8 | -3,8 | -3,8 | -0,7 | -0,7 | -0,7 | -23,6 | -23,6 | -23,6 |
| Bulgaria | 0,7 | 0,7 | 0,7 | -2,6 | -2,6 | -2,6 | -0,2 | -0,2 | -0,2 | -23,3 | -23,3 | -23,3 |
| Spain | 1,3 | 1,3 | 1,3 | 1,3 | 1,3 | 1,3 | -2,3 | -2,3 | -2,3 | -15,6 | -15,6 | -15,6 |
| Slovenia | -0,2 | -0,2 | -0,2 | -1,6 | -1,6 | -1,6 | 1,9 | 1,9 | 1,9 | -22,6 | -22,6 | -22,6 |
| Italy | 2,5 | 2,5 | 2,5 | 0,9 | 0,9 | 0,9 | -0,6 | -0,6 | -0,6 | -30,4 | -30,4 | -30,4 |
| Lithuania | 0,0 | 0,0 | 0,0 | -2,7 | -2,7 | -2,7 | 1,0 | 1,0 | 1,0 | -19,5 | -19,5 | -19,5 |
| Poland | 0,7 | 0,7 | 0,7 | -1,0 | -1,0 | -1,0 | -2,5 | -2,5 | -2,5 | -20,9 | -20,9 | -20,9 |
| UK | -0,2 | -0,2 | -0,2 | -0,9 | -0,9 | -0,9 | 0,7 | 0,7 | 0,7 | -25,8 | -25,8 | -25,8 |

Table-3. Coverage Rates CR (%) of confidence interval for the headcount index P, and using a sampling fraction of $f=1\%$ ($=n/N$)

| Country | \hat{P}_{SRSWOR} | | | \hat{P}_r | | | \hat{P}_{SSRS} | | | $\hat{P}_{P.ST}$ | | |
|-----------|--------------------|-----------------|-------------|----------------|----------------|-----------------|------------------|----------------|----------------|------------------|-------------|----------------|
| | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} |
| Belgium | 92,7 | 92,7 | 92,7 | 93,1 | 93,1 | 93,1 | 93,4 | 93,4 | 93,4 | 90,7 | 90,7 | 90,7 |
| Bulgaria | 94,1 | 94,1 | 94,1 | 93,8 | 93,8 | 93,8 | 94,4 | 94,4 | 94,4 | 90,4 | 90,4 | 90,4 |
| Spain | 94,6 | 94,6 | 94,6 | 94,9 | 94,9 | 94,9 | 94,7 | 94,7 | 94,7 | 93,0 | 93,0 | 93,0 |
| Slovenia | 93,6 | 93,6 | 93,6 | 93,9 | 93,9 | 93,9 | 95,1 | 95,1 | 95,1 | 90,4 | 90,4 | 90,4 |
| Italy | 94,6 | 94,6 | 94,6 | 94,8 | 94,8 | 94,8 | 94,8 | 94,8 | 94,8 | 89,3 | 89,3 | 89,3 |
| Lithuania | 94,5 | 94,5 | 94,5 | 93,2 | 93,2 | 93,2 | 93,8 | 93,8 | 93,8 | 90,0 | 90,0 | 90,0 |
| Poland | 94,3 | 94,3 | 94,3 | 94,6 | 94,6 | 94,6 | 94,7 | 94,7 | 94,7 | 91,7 | 91,7 | 91,7 |
| UK | 94,3 | 94,3 | 94,3 | 94,0 | 94,0 | 94,0 | 94,7 | 94,7 | 94,7 | 90,3 | 90,3 | 90,3 |

Table-4. Percentage of times that the headcount index P is below the lower limits of confidence intervals for the headcount index P, and using a sampling fraction of $f=1\%$ ($=n/N$)

| Country | \hat{P}_{SRSWOR} | | | \hat{P}_r | | | \hat{P}_{SSRS} | | | $\hat{P}_{P.ST}$ | | |
|-----------|--------------------|-----------------|-------------|----------------|----------------|-----------------|------------------|----------------|----------------|------------------|-------------|----------------|
| | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} |
| Belgium | 1,4 | 1,4 | 1,4 | 2,3 | 2,3 | 2,3 | 1,9 | 1,9 | 1,9 | 3,4 | 3,4 | 3,4 |
| Bulgaria | 1,7 | 1,7 | 1,7 | 2,6 | 2,6 | 2,6 | 2,5 | 2,5 | 2,5 | 3,8 | 3,8 | 3,8 |
| Spain | 1,8 | 1,8 | 1,8 | 2,2 | 2,2 | 2,2 | 2,1 | 2,1 | 2,1 | 3,2 | 3,2 | 3,2 |
| Slovenia | 1,4 | 1,4 | 1,4 | 1,9 | 1,9 | 1,9 | 1,7 | 1,7 | 1,7 | 3,1 | 3,1 | 3,1 |
| Italy | 1,8 | 1,8 | 1,8 | 2,0 | 2,0 | 2,0 | 2,2 | 2,2 | 2,2 | 4,2 | 4,2 | 4,2 |
| Lithuania | 1,9 | 1,9 | 1,9 | 2,4 | 2,4 | 2,4 | 1,9 | 1,9 | 1,9 | 3,2 | 3,2 | 3,2 |
| Poland | 1,7 | 1,7 | 1,7 | 2,3 | 2,3 | 2,3 | 2,1 | 2,1 | 2,1 | 3,7 | 3,7 | 3,7 |
| UK | 1,8 | 1,8 | 1,8 | 2,2 | 2,2 | 2,2 | 2,1 | 2,1 | 2,1 | 3,9 | 3,9 | 3,9 |

Table-5. Percentage of times that the headcount index P is above the upper limits of confidence intervals for the headcount index P, and using a sampling fraction of $f=1\%$ ($=n/N$)

| Country | \hat{P}_{SRSWOR} | | | \hat{P}_r | | | \hat{P}_{SSRS} | | | $\hat{P}_{P.ST}$ | | |
|-----------|--------------------|-----------------|-------------|----------------|----------------|-----------------|------------------|----------------|----------------|------------------|-------------|----------------|
| | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} | \hat{V}_{HT} | \hat{V}_{SYG} | \hat{V}_H | \hat{V}_{HT} |
| Belgium | 5,9 | 5,9 | 5,9 | 4,6 | 4,6 | 4,6 | 4,7 | 4,7 | 4,7 | 5,9 | 5,9 | 5,9 |
| Bulgaria | 4,2 | 4,2 | 4,2 | 3,6 | 3,6 | 3,6 | 3,1 | 3,1 | 3,1 | 5,8 | 5,8 | 5,8 |
| Spain | 3,6 | 3,6 | 3,6 | 2,9 | 2,9 | 2,9 | 3,2 | 3,2 | 3,2 | 3,8 | 3,8 | 3,8 |
| Slovenia | 5,0 | 5,0 | 5,0 | 4,2 | 4,2 | 4,2 | 3,2 | 3,2 | 3,2 | 6,5 | 6,5 | 6,5 |
| Italy | 3,6 | 3,6 | 3,6 | 3,2 | 3,2 | 3,2 | 3,0 | 3,0 | 3,0 | 6,5 | 6,5 | 6,5 |
| Lithuania | 3,6 | 3,6 | 3,6 | 4,4 | 4,4 | 4,4 | 4,3 | 4,3 | 4,3 | 6,8 | 6,8 | 6,8 |
| Poland | 4,0 | 4,0 | 4,0 | 3,1 | 3,1 | 3,1 | 3,2 | 3,2 | 3,2 | 4,6 | 4,6 | 4,6 |
| UK | 3,9 | 3,9 | 3,9 | 3,8 | 3,8 | 3,8 | 3,2 | 3,2 | 3,2 | 5,8 | 5,8 | 5,8 |

5. Discussion and Future Research

The analysis and measurement of the poverty is an important topic, and which receives quite attention from many agencies and institution. One of the most important indicators is the headcount index, which is the poverty indicator analyzed in this work.

After analysing numerically the bias and the efficiency of estimators of the headcount index when samples are selected under different sampling designs, we can conclude that the sampling design with better properties is stratified random sampling, i.e., this sampling design gives better results in terms of Relative Bias (RB) and Relative Root Mean Square Error (RRMSE). The second sampling design with better results is simple random sampling with auxiliary information.

In addition, we also evaluated different variance expressions under the same sampling designs. We used the empirical coverage rates of the corresponding confidence intervals and the relative bias of such variance estimators. Our findings are that the various confidence intervals give desirable results, with reasonable coverage rates, excepts

the post-stratified sampling design, which provides variance estimators with large biases and confidence intervals with coverage rate close to 90%.

A further step from this study could consider alternative populations, or include additional sampling designs, such as systematic sampling, cluster sampling, etc. Also the simulation studies could be based on another data basis instead of the 2011 EU-SILC, used in this paper. In addition, this paper focuses on the headcount index. We could extend this study to alternative poverty measures, such as the poverty gap index, the poverty severity index, etc. In particular, it could be interesting to analyze the effect of different sampling designs on multi-dimensional poverty indicators.

In summary, the results derived from this paper and using different sampling designs indicate the advantages of using stratified random sampling when estimating the headcount index. Different variance estimators are also analyzed, and desirable results are also obtained. As a much more general vision, the paper also summarizes the importance of the poverty measurement and the use of statistical instruments as a key factor in the fight against poverty.

Acknowledgement

«This research has been partially supported by the Ministry of Economy, Industry and Competitiveness, the Spanish State Research Agency (SRA) and European Regional Development Fund (ERDF) (project reference ECO2017- 86822-R).»

References

- Álvarez, E., García-Fernández, R. M. and Muñoz, J. F. (2014). A new method to estimate the low income proportion: Monte carlo simulations. *World academy of Science, engineering and Technology. International Journal of Social, Behavioral, Educational, Economic, Business and Industrial Engineering*, 8(10): 3273-76.
- Atkinson, A. B. (1987). On the measurement of poverty. *Econometrica*, 55(4): 749-64.
- Berger, Y. G. and Torres, O. D. L. R. (2014). *Empirical likelihood confidence intervals: an application to the EU-SILC household surveys*. In *Contributions to Sampling Statistics*. Springer, Cham. 65-84.
- Berger, Y. G. and Munoz, J. F. (2015). On estimating quantiles using auxiliary information. *Journal of Official Statistics*, 31(1): 101-19.
- Breidt, F. J. and Opsomer, J. D. (2000). Local polynomial regression estimators in survey sampling. *The Annals of Statistics*, 28(4): 1026-53.
- Chambers, R. L. and Dunstan, R. (1986). Estimating distribution functions from survey data. *Biometrika*, 73(3): 597-604.
- Chen, J. and Sitter, R. (1999). A pseudo empirical likelihood approach to the effective use of auxiliary information in complex surveys. *Statistica Sinica*, 9(2): 385-406.
- Cochran, W. G. (1977). *Sampling techniques*. 3rd edn John Wiley and Sons. 89-146.
- Deville, J. C. and Särndal, C. E. (1992). Calibration estimators in survey sampling. *Journal of the American statistical Association*, 87(418): 376-82.
- Dorfman, A. H. (2009). *Inference on distribution functions and quantiles*. In *Handbook of Statistics 29B Sample surveys: Inference and Analysis*, D. Pfeffermann and C. R. Rao. North-Holland. 371-95.
- Eurostat (2010). *ESS Guidelines for the implementation of the ESS Quality and Performance Indicators*. Office for Official Publications of the European Communities: Luxembourg.
- Eurostat (2013). *Handbook on precision requirements and variance estimation for ESS household surveys*. Publications Office of the European Union: Luxembourg.
- Foster (1998). Absolute versus relative poverty. *The American Economic Review*, 88(2): 335-41.
- Foster, Greer, J. and Thorbecke, E. (1984). A class of decomposable poverty measures. *Econometrica*, 52(3): 761-66.
- Goedemé, T. (2010). The standard error of estimates based on EU-SILC: An exploration through the Europe 2020 poverty indicators. Working Paper No. 10/09. University of Antwerp.
- Goedemé, T. (2013). How much confidence can we have in EU-SILC? Complex sample designs and the standard error of the Europe 2020 poverty indicators. *Social Indicators Research*, 110(1): 89-110.
- Hájek, J. (1964). Asymptotic theory of rejective sampling with varying probabilities from a finite population. *The Annals of Mathematical Statistics*, 35(4): 1491-523.
- Harms, T. and Duchesne, P. (2006). On calibration estimation for quantiles. *Survey Methodology*, 32(1): 37.
- Haughton, J. and Khandker, S. R. (2005). *Introduction to poverty analysis*. World Bank Institute.
- Haughton, J. and Khandker, S. R. (2009). *Handbook on poverty and inequality*. World Bank Publications.
- Horvitz, D. G. and Thompson, D. J. (1952). A generalization of sampling without replacement from a finite universe. *Journal of the American statistical Association*, 47(260): 663-85.
- Khandker, S. R. (2005). *Introduction to poverty analysis*. World Bank Institute.
- López-Escobar, E., Barrios-Zamudio, E. and Muñoz, J. F. (2016). Package *samplingvarest*. The r project for statistical computing.
- Muñoz, J. F., Álvarez-Verdejo, E., García-Fernández, R. M. and Barroso, L. J. (2015). Efficient estimation of the Headcount Index. *Social Indicators Research*, 123(3): 713-32.

- Museux, J. M. (2005). *EU-SILC anonymisation: Results of the eurostat task force*. Monograph of Official Statistics. 153-63.
- Narain, R. D. (1951). On sampling without replacement with varying probabilities. *Journal of the Indian Society of Agricultural Statistics*, 3: 169-74.
- Rao, J. N. K., Kovar, J. and Mantel, H. (1990). On estimating distribution functions and quantiles from survey data using auxiliary information. *Biometrika*, 77(2): 365-75.
- Rao, J. N. K., Wu, C. F. J. and Yue, K. (1992). Some recent work on resampling methods for complex surveys. *Survey Methodology*, 18(2): 209-17.
- Ravallion, M. (1999). *Issues in measuring and modeling poverty*. The World Bank.
- Rueda, M. and Muñoz, J. F. (2011). Estimation of poverty measures with auxiliary information in sample surveys. *Quality and Quantity*, 45(3): 687-700.
- Särndal, C. E., Swensson, B. and Wretman, J. (2003). *Model assisted survey sampling*. Springer Science and Business Media.
- Sen (1953). On the estimate of the variance in sampling with varying probabilities. *Journal of the Indian Society of Agricultural Statistics*, 5(1194): 127.
- Sen (1976). Poverty: an ordinal approach to measurement. *Econometrica*, 44(2): 219-31.
- Silva, P. N. and Skinner, C. J. (1995). Estimating distribution functions with auxiliary information using poststratification. *Journal of Official Statistics*, 11(3): 277.
- Thompson, B. S. (2013). Empirical likelihood-based inference for poverty measures with relative poverty lines. *Econometric Reviews*, 32(4): 513-23.
- Valliant, R. (1993). Poststratification and conditional variance estimation. *Journal of the American Statistical Association*, 88(421): 89-96.
- Van Kerm, P. (2007). Extreme incomes and the estimation of poverty and inequality indicators from EU-SILC. IRISS Working Paper Series 2007-01, CEPS/INSTEAD.
- Wu, C. (1999). *The effective use of complete auxiliary information from survey data*. Simon Fraser University: Burnaby, Canada.
- Yates, F. and Grundy, P. M. (1953). Selection without replacement from within strata with probability proportional to size. *Journal of the Royal Statistical Society: Series B (Methodological)*, 15(2): 253-61.
- Zheng, B. (2001). Statistical inference for poverty measures with relative poverty lines. *Journal of Econometrics*, 101(2): 337-56.